

# MAXFORCE

## GRANULAR FLY BAIT

**FOR PEST MANAGEMENT PROFESSIONALS, LIVESTOCK PRODUCERS, AND COMMERCIAL USE ONLY**

### ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N nitro-2-imidazolidinimine . . . . . 0.50%  
Z-9-Tricosene . . . . . 0.10%

### OTHER INGREDIENTS:

TOTAL . . . . . 100.00%

**STOP - Read the label before each use.  
KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.

#### FIRST AID

IF SWALLOWED:	-Call a poison control center or doctor immediately for treatment advice. -Have person sip a glass of water if able to swallow. -Do not induce vomiting unless told to by a poison control center or doctor. -Do not give anything to an unconscious person.
IF ON SKIN:	-Take off contaminated clothing. -Rinse skin immediately with plenty of water for 15-20 minutes. -Call a poison control center or doctor for treatment advice.
IF INHALED:	-Move person to fresh air. -If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. -Call a poison control center or doctor for further treatment advice.
IF IN EYES:	-Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. -Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. The Emergency Response Telephone No. is 800-334-7577. **Note To Physician:** No specific antidote is available. Treat patient symptomatically.

#### ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates.

Do not apply directly to water, or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. The chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

#### PRO FACTS

- Fast acting
- Ready-to-use scatter bait or paint-on application.
- Contains a specialty formulated mixture of attractants
- Controls flies resistant to organophosphates and carbamates

MAXFORCE GRANULAR FLY BAIT	• Ready-to-use bait containing 0.5% imidacloprid • Contains the attractant Muscalure® • Contains Bitrex® bittering agent to help prevent ingestion by animals, children and pets
CONTROL	• Flies
WHERE TO APPLY	• Around the outside of structures or partially enclosed or protected areas • In and around agricultural production facilities • Around the outside of commercial facilities
MIXING INFORMATION	• Ready-to-use scatter bait • Paint-on application: Mix the fly bait with water
HOW TO APPLY	• Scatter application • Place in any fly bait station • Paint-on application
COVERAGE	• XX Square feet (5.7 to 6.3 oz/1000 sq ft)
RE-APPLY	• Re-apply every 7 days as needed.
QUESTIONS??	• For questions or comments, call toll-free 1-800-843-1702 or visit us online at <a href="http://www.MAXFORCE.com">www.MAXFORCE.com</a>

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **IMPORTANT:** Read entire label and Conditions of Sale before using this product.

EPA Reg. No. 432-1375

EPA Est. No. 7319-NLD001

**NET CONTENTS: 5 or 40 Pounds**

#### FLY MANAGEMENT

WHERE TO APPLY	• For use only around the outside of commercial facilities like canneries, dairy, meat and poultry processing plants, beverage plants, food processing plants, restaurants, bakeries, supermarkets, warehouses, dumpsters, grease pits, recycling areas, loading docks. • For use only outside of agricultural production facilities like broiler houses, feedlots, livestock housing structures, and horse stables. • For use only around the outside of and in walkways of caged layer houses. Place bait in areas inaccessible to animals. • In horse stables, apply bait only in fly bait stations. Place bait stations in areas inaccessible to animals.
----------------	--

USE RATE	• 5.7 to 6.3 oz/1000 sq. ft. (scatter bait and bait station)
HOW TO APPLY	Apply the bait in many locations. Avoid cool or windy locations because flies prefer warm sites for resting. Time applications to begin at the start of the season before fly populations have reached their peak. • <b>Scatter Application:</b> Scatter the bait directly from the directly container onto dry level surfaces so that the individual granules lie near each other without forming small piles. • <b>Bait stations:</b> Place in any fly bait station. Use 1 bait station to cover approximately 250 sq ft and add up to 1.6 oz of bait per station. Secure bait stations at least 4 feet above the ground. Bait stations should be inaccessible to food producing animals, children and pets. • <b>Paint-on application:</b> Mix 1.5oz bait with 1 fl oz warm water (3 lbs bait in 1 quart water) and stir thoroughly. Let stand for about 15 minutes until a paste consistency suitable for painting has formed. Apply the paste with a brush to surfaces where flies rest avoiding surfaces that are dusty.

RESTRICTIONS	• Do not place bait in areas accessible to food producing animals, children and pets. • Do not allow bait to come into contact with food or water. • Do not apply to milking equipment. • Do not apply inside food processing plants, restaurants or other commercial facilities.
--------------	--

#### MANAGEMENT TIPS

- For best results: use the bait as part of an overall integrated Pest Management program.
- Use proper sanitation and manure management.
- Use along with dilutable sprays like TEMPO.
- Use environmental controls, such as proper light management.
- Consult with cooperative Extension.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open containers in a manner as to prevent spillage.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty container. Dispose of container at an approved waste disposal facility or by incineration if allowed by state and local authorities. Do not use container in connection with food, feed or drinking water.  
**SPILLS:** If the container is leaking or material is spilled, carefully sweep up spilled bait and wash exposed area. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Emergency Response Telephone No. is 800-334-7577 or contact Chemtrec at 800-424-9300.

#### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Bayer/Environmental Science's control, it is impossible for Bayer/Environmental Science to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER ENVIRONMENTAL SCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED OF MERCHANTABILITY OR OF FITNESS FOR PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer/Environmental Science is authorized to make any warranties beyond those contained herein. Bayer/Environmental Science disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

**LIMITATION OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Bayer AG, 2005



Bayer Environmental Science  
A Business Group of Bayer CropScience LP

Bayer Environmental Science  
2 T. W. Alexander Drive  
Research Triangle Park, NC 27709  
[www.bayerprocentral.com](http://www.bayerprocentral.com)

**PRODUCT OF HOLLAND**